

# RHEUMATOLOGY ENROLLMENT FORM - "SPECIALTY"

## PATIENT INFORMATION

## TREATMENT ARRANGEMENTS

Last Name \_\_\_\_\_ First Name \_\_\_\_\_  
 Social Security # \_\_\_\_\_ Date of Birth \_\_\_\_\_  
 Address: \_\_\_\_\_  
 Sex  M  F Wgt: \_\_\_\_\_ Ht: \_\_\_\_\_ Phone: \_\_\_\_\_  
 Allergies: \_\_\_\_\_ Cell: \_\_\_\_\_

Start Date: \_\_\_\_\_  
 Ship Meds:  Home  Doctor's Office  
 Teaching by:  Doctor's Office  
 Other: \_\_\_\_\_

## INSURANCE INFORMATION

Insurance:  Medicare  Medicaid  Commercial: \_\_\_\_\_  
 ID \_\_\_\_\_ BIN \_\_\_\_\_ PCN \_\_\_\_\_ GROUP \_\_\_\_\_

**Diagnosis:**  Rheumatoid Arthritis  Ankylosing Spondylitis  Osteoporosis  Psoriatic Arthritis  Lupus  Gout  Spondyloarthropathy  
 Uveitis Date of Diagnosis: \_\_\_\_\_  
**Prior Treatment:**  Humira  Enbrel  Stelara  Others: \_\_\_\_\_ **TB Test**  Positive  Negative

## PRESCRIPTION INFORMATION

MEDICATION	STRENGTH	DOSAGE FORM	DIRECTIONS	QUANTITY	REFILLS
<input type="checkbox"/> Actemra	<input type="checkbox"/> 162mg <input type="checkbox"/> _____mg	<b>INJECTION</b>	<input type="checkbox"/> Less than 100kg weight: Inject 162mg subq every other week, followed by an increase to every week based on clinical response. <input type="checkbox"/> Above 100kg weight: Inject 162mg subq every week. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Cimzia	<input type="checkbox"/> 200mg/1ml Prefilled Syringe <input type="checkbox"/> 200mg Vial		<input type="checkbox"/> <b>Induction Dose:</b> Inject 400mg subcutaneously on day 1, at week 2, and at week 4. <input type="checkbox"/> <b>Maint. Dose:</b> Inject 200mg subcutaneously every other week. <input type="checkbox"/> <b>Maint. Dose:</b> Inject 400mg subcutaneously every 4 weeks.		
<input type="checkbox"/> Cosentyx	<input type="checkbox"/> 300mg <input type="checkbox"/> Pen <input type="checkbox"/> PFS <input type="checkbox"/> 150mg <input type="checkbox"/> Pen <input type="checkbox"/> PFS		<input type="checkbox"/> <b>Initial Dose:</b> Inject 300mg subq at weeks 0,1,2,3, and 4 weeks. Each 300mg dose is given as 2 subq inj of 150mg. <input type="checkbox"/> <b>Maint Dose:</b> Inject 300mg subq once 4 weeks. Each 300mg dose is given as 2 subq inj of 150mg.		
<input type="checkbox"/> Enbrel	<input type="checkbox"/> SureClick® Pen <input type="checkbox"/> Mini™ with Auto Touch <input type="checkbox"/> Prefilled Syringe <input type="checkbox"/> 25mg <input type="checkbox"/> 50mg <input type="checkbox"/> Vials 25mg		<input type="checkbox"/> Inject 50mg SQ every week <input type="checkbox"/> Inject 25mg SQ TWICE a week (72-96 hours apart). <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Humira	<input type="checkbox"/> Uveitis Starter Kit <input type="checkbox"/> 40mg Pen <input type="checkbox"/> 40mg Pre-Filled Syringe <input type="checkbox"/> 80mg Pen		<input type="checkbox"/> <b>Uveitis Initial Dose:</b> Inject 80mg Subq on day 1, 40mg on day 8 and 40mg day 22 then maintenance. <input type="checkbox"/> <b>Maintenance Dose:</b> Inject 40mg Subq every other week. <input type="checkbox"/> <b>Maintenance Dose:</b> Inject 40mg Subq weekly.		
<input type="checkbox"/> Humira CF (Citrate-free)	<input type="checkbox"/> Uveitis Starter Kit <input type="checkbox"/> 40mg Pen <input type="checkbox"/> 40mg Pre-Filled Syringe <input type="checkbox"/> 80mg Pen		<input type="checkbox"/> <b>Uveitis Initial Dose:</b> Inject 80mg Subq on day 1, 40mg on day 8 and 40mg day 22 then maintenance. <input type="checkbox"/> <b>Maintenance Dose:</b> Inject 40mg Subq every other week. <input type="checkbox"/> <b>Maintenance Dose:</b> Inject 40mg Subq weekly.		
<input type="checkbox"/> Kevzara	<input type="checkbox"/> 200mg <input type="checkbox"/> 150mg		<input type="checkbox"/> Inject 200mg subcutaneously once every two weeks. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Otrexup	<input type="checkbox"/> _____mg/ml		<input type="checkbox"/> Inject _____mg Subcutaneously Once a week.		
<input type="checkbox"/> Rasuvo	<input type="checkbox"/> _____mg/ml		<input type="checkbox"/> Inject _____mg Subcutaneously Once a week.		
<input type="checkbox"/> Stelara	<input type="checkbox"/> 45mg Prefilled Syringe <input type="checkbox"/> 90mg Prefilled Syringe		<input type="checkbox"/> Inject 45mg SQ at weeks 0, 4 & then every 12 weeks. <input type="checkbox"/> Inject 90mg SQ at weeks 0, 4 & then every 12 weeks.		
<input type="checkbox"/> Simponi	<input type="checkbox"/> 50mg/0.5ml Prefilled SmartJect Autoinjector <input type="checkbox"/> 50mg/0.5ml Prefilled Syringe		<input type="checkbox"/> Inject 50mg (0.5ml) subcutaneously once a month. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Forteo	<input type="checkbox"/> 600ug/2.4 ml Delivery Device		<input type="checkbox"/> Inject 20ug (0.08 ml) Subcutaneously Once daily.		
<input type="checkbox"/> Prolia	<input type="checkbox"/> 60mg/ml		<input type="checkbox"/> Inject 60mg/ml Subcutaneously Once every 6 months. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Tymlos	<input type="checkbox"/> 3120mcg/1.56ml		<input type="checkbox"/> Inject 80mcg Subq once daily.		
<input type="checkbox"/> Taltz	<input type="checkbox"/> 80mg/ml Autoinjector <input type="checkbox"/> 80mg/ml Prefilled Syringe	<input type="checkbox"/> Inject 160mg Subcutaneously (two 80mg injections) at week 0, followed by 80mg every 4 weeks. <input type="checkbox"/> Other: _____			
<input type="checkbox"/> Otezla	<input type="checkbox"/> Starter Pack <input type="checkbox"/> 30mg Tablet	<b>ORAL</b>	<input type="checkbox"/> Titrate: Take 1 tablet on day 1 then twice daily as directed <b>OR date provided</b> _____ <input type="checkbox"/> Maintenance: Take 1 tablet by mouth twice daily. <input type="checkbox"/> Bridge RX: Take 1 tablet by mouth twice daily; disp. by OSP.		
<input type="checkbox"/> Olumiant	<input type="checkbox"/> 2mg		<input type="checkbox"/> Take 2mg PO Once daily.		
<input type="checkbox"/> Xeljanz	<input type="checkbox"/> 5mg		<input type="checkbox"/> Take 5mg PO Twice a day.		
<input type="checkbox"/> Xeljanz XR	<input type="checkbox"/> 11mg		<input type="checkbox"/> Take 11mg PO Once daily.		

Physician Signature: \_\_\_\_\_  DAW (Dispense as Written) Date: \_\_\_\_\_  
 Physician Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Office Contact: \_\_\_\_\_  
 Physician Address: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_

IMPORTANT NOTICE: This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake. Then destroy this document. Please direct all verification or notification to the above mentioned physician's office.  
 X By Signing this prescription & using Southside's services, you are authorizing Southside to contact insurance companies for prior authorizations on behalf of your practice.



**FAX TO (855) - 822 - 7838**  
 Specialty Pharmacy | IV Therapy | Home Health | Medical Equipment | Physician's Billing  
 7700 Main St., Houston, TX 77030 | P: 855-822-7828 | www.ssr.com

**"One SOURCE for ALL your Medical NEEDS"**

# RHEUMATOLOGY ENROLLMENT FORM - "INFUSION"

## PATIENT INFORMATION

Last Name \_\_\_\_\_ First Name \_\_\_\_\_  
 Social Security # \_\_\_\_\_ Date of Birth \_\_\_\_\_  
 Address: \_\_\_\_\_  
 Sex  M  F Wgt: \_\_\_\_\_ Ht: \_\_\_\_\_ Phone: \_\_\_\_\_  
 Allergies: \_\_\_\_\_ Cell: \_\_\_\_\_

## TREATMENT ARRANGEMENTS

Start Date: \_\_\_\_\_  
 Ship Meds:  Home  Doctor's Office  
 Teaching by:  Doctor's Office  
 Other: \_\_\_\_\_

**Diagnosis:**  Rheumatoid Arthritis  Ankylosing Spondylitis  Osteoporosis  Psoriatic Arthritis  Lupus  Gout  Spondyloarthropathy  
 Date of Diagnosis: \_\_\_\_\_

## INSURANCE INFORMATION

**Insurance:**  Medicare  Medicaid  Commercial: \_\_\_\_\_  
 ID \_\_\_\_\_ BIN \_\_\_\_\_ PCN \_\_\_\_\_ GROUP \_\_\_\_\_

**Premeds:**  Tylenol 500mg 2 PO  Loratidine 10mg PO  Zyrtec 10mg PO  Benadryl \_\_\_\_\_ mg IV/PO (circle one)  
 SoluCortef \_\_\_\_\_ mg IV  Ondansetron \_\_\_\_\_ mg IV  Promethazine \_\_\_\_\_ mg IV  Other \_\_\_\_\_

Standing Order: Anaphylaxis Protocol  **Skilled Nurse to start PIV, infuse per protocol, DC PIV each visit**

Lab Draw As Follows: \_\_\_\_\_

Quantiferon Gold Lab Draw Q \_\_\_\_\_  Patient to FU w/MD Q \_\_\_\_\_

**Prior Treatment:**  Humira  Enbrel  Stelara  Others: \_\_\_\_\_ **TB Test**  Positive  Negative

## PRESCRIPTION INFORMATION

MEDICATION	STRENGTH	DOSAGE FORM	DIRECTIONS	QUANTITY	REFILLS
<input type="checkbox"/> Actemra	<input type="checkbox"/> 80mg/4ml <input type="checkbox"/> 200mg/10ml <input type="checkbox"/> 400mg/20ml <input type="checkbox"/> _____ mg/kg	IV	<input type="checkbox"/> <b>Induction Dose:</b> 4 mg/kg every 4 weeks. <input type="checkbox"/> <b>Maint. Dose:</b> (based on clinical response): 8 mg/kg every 4 weeks. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Benlysta	<input type="checkbox"/> 10mg/kg		<input type="checkbox"/> <b>Induction Dose:</b> Administer 10 mg/kg on week 0, week 2 week 4. <input type="checkbox"/> <b>Maint. Dose:</b> 10 mg/kg once every 4 weeks after Induction dose. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Boniva	<input type="checkbox"/> 3mg/3ml Injection		<input type="checkbox"/> Inject 3mg/ml Once every 3 months. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Krystexxa	<input type="checkbox"/> 8mg/ml		<input type="checkbox"/> Inject 8mg/ml Intravenous once every 2 weeks. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Orencia	<input type="checkbox"/> 250mg Vial <input type="checkbox"/> _____ mg/kg		<input type="checkbox"/> Infuse _____ mg in 100ml of 0.9% NaCl at weeks 0, 2, and 4, then every 4 weeks thereafter. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Remicade	<input type="checkbox"/> 100mg Vial <input type="checkbox"/> 300mg/kg		<input type="checkbox"/> <b>Induction Dose:</b> IV in 250ml of 0.9% NaCl at weeks 0, 2, and 6 weeks. <input type="checkbox"/> <b>Maint. Dose:</b> IV in 250ml of 0.9% NaCl every 8 weeks. <input type="checkbox"/> <b>Maint. Dose:</b> IV in 250ml of 0.9% NaCl every 6 weeks. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Rituxan	<input type="checkbox"/> 100mg/10ml vial <input type="checkbox"/> 500mg/50ml vial		<input type="checkbox"/> Infuse two doses of 100mg in 1 liter of 0.9% NaCl separated by 2 weeks. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Reclast	<input type="checkbox"/> 5mg/100ml		<input type="checkbox"/> 5mg/100ml Once every year. <input type="checkbox"/> Other: _____		
<input type="checkbox"/> Simponi Aria	<input type="checkbox"/> _____ mg/kg		<input type="checkbox"/> Inject 2 mg/kg intravenous (IV) infusion over 30 minutes at Weeks 0 and 4, then every 8 weeks thereafter <input type="checkbox"/> Other: _____		

Physician Signature: \_\_\_\_\_  DAW (Dispense as Written) Date: \_\_\_\_\_

Physician Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Office Contact: \_\_\_\_\_

Physician Address: \_\_\_\_\_ NPI: \_\_\_\_\_ DEA: \_\_\_\_\_

**IMPORTANT NOTICE:** This message may contain privileged and confidential information and is intended only for the individual named. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please notify the sender immediately if you have received this document by mistake. Then destroy this document. Please direct all verification or notification to the above mentioned physician's office.

X By Signing this prescription & using Southside's services, you are authorizing Southside to contact insurance companies for prior authorizations on behalf of your practice.



Specialty Pharmacy | Infusion Services

**FAXTO (855) - 822 - 7838**

Specialty Pharmacy | IV Therapy | Home Health | Medical Equipment | Physician's Billing

7700 Main St., Houston, TX 77030 | P: 855-822-7828 | www.ssr.com

**"One SOURCE for ALL your Medical NEEDS"**